

**AN INTEGRAL OF CHOI KWOK PUI APPEARING IN A
POISSON PROCESS APPROXIMATION**

In the calculation of an immigration-death process, close to a Poisson process, it was needed to evaluate the expectation

$$\int_0^{\infty} e^{-t} E \left(\frac{1}{Y(t) + Z(t) + 1} \right) dt$$

where $Y(t)$ and $Z(t)$ are independent, $Y(t)$ is distributed as Binomial(m, e^{-t}) and $Z(t)$ is Poisson distributed with mean $\lambda(1 - e^{-t})$. This leads to the evaluation of the double integral

$$(1) \quad F(\lambda, m) = \int_0^1 \int_0^1 (1 - r + ur)^m e^{-\lambda ur} du dr.$$

Here $m \in \mathbb{N}$ and $\lambda \in \mathbb{R}^+$.

The integral is given by

$$(2) \quad F(\lambda, m) = \frac{m!}{\lambda^{m+1}} \left[\left(\sum_{k=0}^m \frac{\lambda^k}{k!} \right) \times \int_0^1 \frac{1 - e^{-\lambda r}}{r} dr + \sum_{k=0}^{m-1} \frac{\lambda^k}{k!} \sum_{\nu=1}^{m-k} \frac{(-\lambda)^\nu}{\nu \nu!} \right].$$

The integral appearing in (2) is evaluated by Mathematica as

$$(3) \quad \int_0^1 \frac{1 - e^{-\lambda r}}{r} dr = \gamma + \ln \lambda + \Gamma(0, \lambda)$$

where γ is the Euler constant

$$(4) \quad \gamma = \lim_{n \rightarrow \infty} 1 + \frac{1}{2} + \cdots + \frac{1}{n} - \ln n \sim 0.577215$$

and $\Gamma(0, x)$ is the incomplete gamma function defined by

$$(5) \quad \Gamma(0, x) := \int_x^{\infty} \frac{e^{-t}}{t} dt.$$

The sum appearing in the first term in (2) can be expressed as

$$(6) \quad \sum_{k=0}^m \frac{\lambda^k}{k!} = \frac{e^\lambda}{m!} \Gamma(m+1, \lambda)$$

so that

$$F(\lambda, m) = \frac{e^\lambda}{\lambda^{m+1}} \Gamma(m+1, \lambda) [\gamma + \Gamma(0, \lambda) + \ln \lambda] + \frac{m!}{\lambda^{m+1}} \sum_{k=0}^{m-1} \frac{\lambda^k}{k!} \sum_{\nu=1}^{m-k} \frac{(-\lambda)^\nu}{\nu \nu!}.$$

General information about this function can be obtained in [2], Chapter 45.

To evaluate the function $F(\lambda, m)$, let $v = 1 - u$ and expand the term $(1 - rv)^m$ to obtain

$$(7) \quad F(\lambda, m) = \sum_{k=0}^m (-1)^{m-k} \binom{m}{k} \int_0^1 r^{m-k} e^{-\lambda r} \int_0^1 e^{\lambda v r} v^{m-k} dv dr.$$

Now

$$(8) \quad \int_0^1 e^{\lambda v r} v^{m-k} dv = \lambda^{-m-1+k} r^{-m-1+k} \int_0^{\lambda r} x^{m-k} e^x dx.$$

The indefinite integral of $x^{m-k} e^x$ can be obtained by repeated integration by parts and it can be found in [1], 2.321.2:

$$(9) \quad \int x^{m-k} e^x dx = e^x \sum_{j=0}^{m-k} \frac{(-1)^j (m-k)!}{(m-k-j)!} x^{m-k-j}$$

The formula (2) appears directly from here. In the sum (7) one should combine the term corresponding to $k = m$ with the inner term corresponding to $j = m - k$ to produce terms involving the integrand $(1 - e^{-\lambda r})/r$. These are integrable near $r = 0$.

The function $F(\lambda, m)$ can be simplified even further using

$$\begin{aligned} \sum_{k=0}^{m-1} \sum_{j=1}^{m-k} \frac{(-1)^j \lambda^{k+j}}{j j! k!} &= \sum_{j+k \leq m} \frac{(-1)^j \lambda^{k+j}}{j j! k!} \\ &= \sum_{r=1}^m \sum_{j=1}^r \frac{(-1)^j \lambda^r}{j j! (r-j)!} \\ &= \sum_{r=1}^m \frac{\lambda^r}{r!} \sum_{j=1}^r \frac{(-1)^j}{j} \binom{r}{j} \\ &= \sum_{r=1}^m \frac{\lambda^r}{r!} \sum_{j=1}^r \frac{1}{j} \\ &= \sum_{r=1}^m \frac{\lambda^r H_r}{r!} \end{aligned}$$

where $H_r = 1 + 1/2 + \dots + 1/r$ is the harmonic number. Therefore

$$F(\lambda, m) = \frac{e^\lambda}{\lambda^{m+1}} \Gamma(m+1, \lambda) [\gamma + \Gamma(0, \lambda) + \ln \lambda] + \frac{m!}{\lambda^{m+1}} \sum_{r=1}^m \frac{\lambda^r H_r}{r!}.$$

REFERENCES

- [1] I.S. Gradshteyn - I.M. Rhyzik: *Tables of Integrals, Series and Products*, 6th edition, Alan Jeffrey and D. Zwillinger editors, Academic Press, 2000.
- [2] J. Spanier - K. Oldham: *An atlas of functions*. Springer-Verlag, 1987.