

PROOF OF FORMULA 3.365.2

$$\int_u^\infty \frac{xe^{-xt} dx}{\sqrt{x^2 - u^2}} = uK_1(tu)$$

The integral representation for the Bessel function K_ν appears in **8.432.3**:

$$K_\nu(z) = \frac{z^\nu \sqrt{\pi}}{2^\nu \Gamma(\nu + 1/2)} \int_1^\infty e^{-tz} (t^2 - 1)^{\nu-1/2} dt.$$

In particular

$$K_0(z) = \int_1^\infty \frac{e^{-zt} dt}{\sqrt{t^2 - 1}} \quad \text{and} \quad K_1(z) = z \int_1^\infty e^{-zt} \sqrt{t^2 - 1} dt.$$

The change of variables $x = tu$ gives

$$K_0(z) = \int_u^\infty \frac{e^{-zx/u} dx}{\sqrt{x^2 - u^2}}.$$

Therefore

$$\int_u^\infty \frac{xe^{-xz} dx}{\sqrt{x^2 - u^2}} = uK'_0(uz).$$

The relation $K'_0(z) = -K_1(z)$ can be established by integration by parts the identity

$$K'_0(z) = - \int_1^\infty \frac{te^{-zt} dt}{\sqrt{t^2 - 1}}.$$

This gives the result.